

An ELISA for studying Ebola Virus / VLP release

George Kallstrom, M. Javad Aman, Sina Bavari
Sina Bavari Lab, USAMRIID
Ft. Detrick, MD

Opinions, interpretations, conclusions, and recommendations are those of the author and are not necessarily endorsed by the U.S. Army. The research described herein was sponsored by the MBD RP research plan titled MMS-Bases Ebo/Mbg Tx / 03-4-4J-012.

17 Nov 04

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 17 NOV 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE An ELISA for studying Ebola Virus / VLP release				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) USAMRIID Ft. Detrick, MD				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001849, 2004 Scientific Conference on Chemical and Biological Defense Research. Held in Hunt Valley, Maryland on 15-17 November 2004 . , The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 23	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Introduction/Agenda


- Filovirus Background
 - Replication
 - Structure
 - VP40
- ELISA setup and optimization
- Effects of Ebola proteins on VLP release
- Effects of VP40 mutants on VLP release
- Therapeutic compound screening

Filoviruses


- *Mononegavirales, Filoviridae*-2 species in family (Marburg and Ebola)
- Both clinically manifest as a rapidly progressive hemorrhagic fever with high mortality rates
- Marburg was weaponized and stockpiled by the USSR
- Sporadic outbreaks have been on the rise in Africa over the past several years
- Unknown Reservoir

Viral Replication


- Single strand negative sense RNA genome
- 7 genes, single segment ~19,000bp



Filovirus Structure




Longitudinal 2D
Filamentous Particles



Transverse 2D section

VP40-Matrix Protein




*Exists in both monomeric and oligomeric conformations

Dessen et al, 2000 EMBO Journal

Objectives


- Develop an assay that measures Ebola virus / VLP release
- Study the effects of Ebola proteins on VLP production
- Determine the regions of VP40 that are important for viral release
- Screen for potential anti-viral therapeutic compounds able to inhibit viral release

Capture ELISA




Standardizing the assay

Purified VP40 protein



- Assay detects ~2ng of VP40 protein
- Linear from 10-125ng


Inactivated Ebola virus




- Assay detects ~500pfu iEbola.
- Linear from 7000-100,000pfu

Detecting Ebola virus in non-human primate samples

Dilutions of Liver Homogenate



Plasma samples





Ebola virus growth curve


ELISA / plaque assay

ELISA (1day)-BL2

Plaque assay (>1 week)-BL4



Standard Curve




Time Point	Plaque Assay	ELISA
48h	1.2×10^7 pfu/ml	1.0×10^7 pfu/ml
72h	3.4×10^8 pfu/ml	1.7×10^8 pfu/ml
96h	2.7×10^8 pfu/ml	2.4×10^8 pfu/ml

BL4 work, D. Swenson


VLP Structure, GP+VP40

- VP40 alone makes filamentous structures that resemble native virus.
- Adding GP increases the production of VLPs and makes the VLPs immunogenic.

Ebola virus (F. Murphy, CDC 1976)




GP+VP40 Virus Like Particle



Standard VLP assay conditions

Transfect
VLP plasmids
into 6 well plate
@80% confluence




Incubate 48h


Harvest cell culture supernatant
Wash the cells with PBS, harvest wash
Harvest cell pellet and disrupt cells for cellular lysates

Perform ELISA

Testing VLP Supernatants / Lysates




Can detect protein in 2ul of supernatant,
linear range 12.5-75ul



Can detect protein in <1ul of lysate,
linear range 3-25ul

Effect of other plasmids on VLP formation



VP40	+	+	+	+	+	+	+	+	+	+	+
GP	-	+	-	-	-	-	+	+	+	+	+
NP	-	-	+	-	-	-	+	+	+	+	+
VP35	-	-	-	-	+	-	-	-	+	+	+
VP30	-	-	-	+	-	-	-	+	-	+	+
VP24	-	-	-	-	-	+	-	-	-	-	+


-GP and NP increase VP40 VLP release

-VP30 slightly inhibits release in the presence of VP40 and NP

-VP35 promotes VLP release in the presence of VP40 and NP

VP40 Oligomerization



Anti-VP40 Western



- VP40 exists in both monomeric and oligomeric forms.
- VP40 oligomers are SDS resistant, but heat sensitive.
- Oligomerization of VP40 is important for formation of VLPs.


VLP/oligomer formation

15amino acid deletions



T. Nelle constructed plasmids


Shaded deleted regions made oligomer / released VLPs



Dessen et al, 2000 EMBO Journal


ELISA can be utilized for therapeutic drug screening

- Identification of compounds that inhibit viral egress.




Potential for therapeutic screening.

Sup from 96 well plate



40,000 cells / well is optimal for splitting/signal, at 50,000 wells are overgrown

VLP release 96 well DMSO treated



<2.5%DMSO did not affect VLP production or cell viability

Accomplishments

- Developed and optimized ELISA that measures Ebola virus / VLP release
- Quantified the contribution of filoviral proteins on VLP formation
- Identified regions of VP40 that are critical for VLP formation release
- Prepared for antiviral therapeutic compound screening

Summary

- We showed a sensitive ELISA for Ebola VLP / viral release
- Although not shown, the ELISA is useful for studying viral-host cell interactions
- Potentially useful as an alternative to plaque assays

Acknowledgements

Sina Bavari Lab

- Sina Bavari-leader of research group
- Javad Aman-research colleague
- Kelly Warfield & Dana Swenson-BSL4 experiments
- Gordon Ruthel-microscopy

Other USAMRIID colleagues

- Shannon Mort-Undergraduate Student
- Sally Charles-Undergraduate Student
- Shirin Badie-Graduate Student
- Tim Nelle-constructed 15aa Δ -VP40 plasmids
- Other lab members and USAMRIID staff who support our research